

PRINTED NAME _____

SCHOOL _____

POSITION _____

DATE COMPLETED _____

Rectal Diazepam – Diastat Acudial® Test

1. What is the definition of status epilepticus?
 - a. Continuous seizure activity lasting 5-10 minutes
 - b. Intermittent seizure activity lasting 30 minutes or more with a return of consciousness
 - c. Diagnosis of person with a seizure disorder
 - d. None of the above

2. Interruption of seizure activity is important for all of the following except:
 - a. Prevention of brain damage
 - b. Prevention of death due to ongoing seizure activity
 - c. Decreasing the frequency of subsequent seizure activity
 - d. Elimination of the need for daily medication to control seizure activity

3. The three 3's refers to counting to three while performing all of the following activities except:
 - a. Delivery of the medication in the rectum
 - b. Before removing syringe from rectum
 - c. Insertion of the syringe in the water soluble lubricant
 - d. Hold the buttock after withdrawal of the syringe

4. If alone with student, what immediate action is recommended after the administration of Diastat®?
 - a. Encourage the student to become physically active
 - b. Feed the student a snack with carbohydrate and protein foods
 - c. Call emergency medical services or 911
 - d. Monitoring the student's gait

5. Before administration of Diastat®, school personnel should make sure the Diastat® has the ready collar exposed on the barrel of the syringe. What color is the ready collar?
 - a. Green
 - b. Brown
 - c. Yellow
 - d. Red