

PRINTED NAME \_\_\_\_\_

SCHOOL \_\_\_\_\_

POSITION \_\_\_\_\_

DATE COMPLETED \_\_\_\_\_

### Epinephrine-EpiPen® or Twinject® Test

1. What other diseases increase the likelihood of future anaphylactic reaction in a particular student?
  - a. Asthma and sickle cell disease
  - b. Sickle cell disease and eczema
  - c. Asthma and eczema
  - d. Asthma, eczema, and sickle cell disease
  
2. Which of the following symptoms is not associated with an anaphylactic reaction?
  - a. Involuntary movements
  - b. Drooling
  - c. Nausea and vomiting
  - d. Tingling of skin
  
3. What percentages of students with first anaphylactic reaction have a second reaction?
  - a. 50%
  - b. 10%
  - c. 85%
  - d. 30%
  
4. How long might a student experience a second anaphylactic reaction after having an initial reaction?
  - a. 24 hours
  - b. 72 hours
  - c. 1 hour
  - d. 16 hours
  
5. What is considered to be the best sequence of steps for administration of epinephrine if alone with the student?
  - a. Administration of the epinephrine, call 911 and parent, monitor for secondary symptoms.
  - b. Call 911 and parent, administer epinephrine, monitor for secondary reactions.
  - c. Administration of the epinephrine, monitor for secondary symptoms, and call 911 and parent.
  - d. None of the options are correct because I do not need to call 911.